

Ascension Island Biodiversity Action Plan

ASCENSION SPURGE



Photo: AIG Conservation

SUMMARY

Taxonomy: Kingdom: Plantae; Phylum: Tracheophyta; Class: Magnoliopsida; Order: Euphorbiales; Family: Euphorbiaceae; Species: *Euphorbia origanoides*

Nativeness: Endemic to Ascension Island

Description: Perennial, dwarf shrub that forms low hemispherical domes up to 1m diameter. Stems are usually reddish and dichotomously branched with clusters of small creamish-white flowers at the tips. Occurs on arid cinder plains and scoria cones at lower elevations.

IUCN Red List status: Critically Endangered 

Local trend: Probably declining 

Threats: The major threats to *E. origanoides* are from introduced competitors and herbivores; secondary threats include climate change and commercial development.




Distribution

Global

Endemic to Ascension Island.

Local

The current distribution of *E. origanoides* is localised and highly fragmented, with most extant sub-populations restricted to a few sparsely-vegetated areas at altitudes between sea level and 150m. Significant population centres are now only found around South Gannet Hill and Cotar Hill in the south west of the Island, the Letterbox Peninsula and Wig Hill in the south east and scattered along the coast from Spire Beach to Hummock Point. Smaller satellite populations comprising of less than 70 individuals are also found in the Comfortless Cove and Sister’s Peak areas. However, *E. origanoides* was probably once widespread in low lying areas and may have had a broader altitudinal range [1]. As recently as 1976 the distribution was more extensive and included populations at North East Bay, Cross Hill and English Bay which have since disappeared [1]. The disappearance of the English Bay population appears to have coincided with the development of the BBC Atlantic Relay Station at this site in 1964 [1].



Current and historic distribution of *Euphorbia origanoides*. Distribution for 1958 and 1976 digitised from [2,3]. Current distribution taken from March 2014 census; symbol sizes are scaled according to total number of individuals encountered (AIG Conservation, unpublished data).

3. Status

Population estimate:	1889 – 14900 individuals	Trend:	Probably declining	IUCN status:	Critically Endangered
----------------------	--------------------------	--------	--------------------	--------------	-----------------------

Determining population trends for *E. origanoides* is challenging due to the substantial variation in its abundance from year to year. Numbers of plants in known populations have fluctuated between 1889 and 14900 individuals since 2008, primarily as a function of varying amounts of rainfall. Indeed, populations appear to be able to persist for long periods as a dormant seed bank, regenerating rapidly when conditions are favourable. Overall trends may therefore be best inferred from long-term changes in distribution, which suggest a 50 % reduction in the area occupied since the 1950s [1].

4. Ecology

Habitat

E. origanoides is a dwarf shrub found on arid, low lying plains of pyroclastic deposits and scoria cones from sea level to an altitude of approximately 300m [1]. Rainfall in these areas is low and episodic and vegetation is sparse. Native associates include *Aristida ascensionis* (grass), *Cyperus appendiculatus* (sedge) and *Portulaca oleracea* (purslane). Mature plants form low hemispherical domes which may extend over 1 m in diameter but seldom attain heights greater than 40 cm [1]. It is often found in small crevices and gullies which probably have lower evapotranspiration rates and increased water availability compared to more exposed sites.

Reproduction & life history

E. origanoides is perennial and flowers year round. Flowers are monoecious (possess both male and female parts) and are insect pollinated, probably by Diptera (flies) and Hemiptera (bugs) [1]. Seed dispersal is predominantly via wind, although water may be an important dispersal agent following heavy deluges [1]. Seeds remain viable for up to 5 years with germination triggered by moisture [1].

Taxonomy & population structure



Some morphological variation between populations is apparent [1] and there is evidence of genetic divergence between east and west coast populations, potentially linked to a dispersal barrier to seeds and pollinators caused by the prevailing south-easterly trade winds [1]. The closest relative is thought to be *E. trinervia* which is native to coastal regions of tropical West Africa [1].

5. Threats*

8.1.2 Invasive non-native/alien species/diseases (named species)

Impact: HIGH

The rapid spread of invasive vegetation across much of the low-lying, ash plain habitat favoured by Ascension spurge represents a major threat to the survival of this species in the wild and has undoubtedly contributed to a continuum of decline. Although the areas occupied by extant populations remain sparsely vegetated, Mexican thorn (*Prosopis juliflora*), tree tobacco (*Nicotiana glauca*), guava (*Psidium guajava*), *Heliotropium curassavicum*, *Casuarina equesitifolia*, wild tomato (*Solanum* sp.) and *Waltheria indica* all have the potential to out-compete Ascension spurge in its native range [4,5]. Mexican thorn in particular is highly efficient at monopolising available resources and now dominates the vegetation across a large swathe of the coastal lowlands, including some former strongholds of Ascension spurge [1]. Grazing by introduced vertebrates and invertebrate pests pose additional and potentially more immediate threats. Six species of introduced mealy bugs and scale insects (*Phenacoccus solani*, *Ferrisia malvastra*, *Planococcus minor*, *Hemiberlesia lataniae*, *Pinnaspis strachani* and *Icerya purchasi*) and one whitefly (*Bemisia tabaci*) are known to attack Ascension spurge, and are having noticeable impacts on plant vigour and survival [6]. Introduced herbivores, particularly rabbits and mice, have also been implicated in population declines [1,4,5] and may be increasing in number due to release from predation by feral cats following their eradication in 2004 [5,7].

1.2 Commercial & industrial development

Impact: LOW

The largest extant population of Ascension spurge at Mars Bay is situated within an area of relatively frequent development and maintenance works associated with Wideawake Airfield and the US Air Force installations at Cotar Hill and South Gannet Hill. Mitigation studies have been conducted previously to lessen the impacts of unavoidable construction works on the spurge population [8] and it is important that this proactive approach continues.

11.2 Climate change & severe weather: Droughts

Impact: UNKNOWN

In the arid habitats occupied by Ascension spurge, rainfall is the principle driver of seedling recruitment and population dynamics. Mass seedling recruitment occurs following episodic periods of heavy rainfall while prolonged periods of drought are associated with significant population declines [5]. It is difficult to predict long term changes in precipitation at Ascension Island with any degree of confidence [9]; however even small reductions in the frequency or quantity of rainfall could have significant impacts on spurge populations growing in already marginal environments.

11.3 Climate change & severe weather: Temperature extremes

Impact: LOW

Temperature is an important controlling factor of evapotranspiration rates and therefore on plant soil-water relations. According to global climate projections, temperatures on Ascension will increase by several degrees over the next 50-100 years [9], threatening to further reduce available soil moisture and increase the risk of heat stress in the already marginal habitats occupied by Ascension spurge [5,9]. Altitudinal range shifts to track changing climate are unlikely to be possible due to the dense coverage of invasive vegetation at higher elevations.

10.3 Avalanches/landslides

Impact: LOW

Several of the extant populations of Ascension spurge occur either partially or wholly on loose, unstable scree slopes and scoria cones, rendering them vulnerable to landslides. In 2013 a landslip completely destroyed the spurge population on Wig Hill which had formerly accounted for 10-15 % of the global population. As of September 2014, no regeneration had yet occurred suggesting that much of the seed bank contained within the topsoil layer may also have been lost.

8.3 Introduced genetic material

Impact: UNKNOWN

Hybridisation with closely related, introduced species may be a pervasive threat to the *E. origanoides* gene pool. Although hybridisation in the subgenus *Chamaesyce* is rare, plants that appear to be intermediate for certain characters have been recorded at the Airhead and Comfortless Cove areas [4]. Further research is needed to



determine whether hybridisation is occurring.

*Threats are classified and scored according to the [IUCN-CMP Unified Classification of Direct Threats](#) [10]

Relevant policies and legislation

Local

Ascension spurge is protected under the [Wildlife Protection Ordinance 2013](#), which prohibits the damaging, killing or possession of protected species without license.

Populations at Letterbox and South Gannet Hill are wholly or partially contained within nature reserves designated under the [National Protected Areas Order 2014](#). The [National Protected Areas Regulations 2014](#) restrict all forms of development within nature reserves.

Management notes

Management for this species is reasonably well-advanced. Propagation protocols are well established and significant numbers of individuals have been successfully cultivated from seed and restored into fenced, semi-natural restoration sites. Grazing exclosures have also been created around important concentrations of wild plants. These measures have been successful in protecting clusters of mature individuals that can help to seed surrounding areas, but are limited in scale and do not offer habitat-level solutions to the threats faced by *E. origanoides*. Indeed, given the significant natural fluctuations in the population size of this species and its tendency to wait out periods of drought as a persistent seed bank, maintaining areas of suitable habitat for regeneration is probably of higher priority than protecting existing plants or restoring cultivated individuals into existing population centres. Further work should focus on controlling invasive species around remaining strongholds and establishing new population centres within the species former range.

Preventing the spread of invasive shrubs and trees such as *Prosopis juliflora*, *Psidium guajava*, *Nicotiana glauca* and *Casuarina equisetifolia* into the remaining strongholds of *E. origanoides* is particularly critical to the long term survival of the species and should be achievable provided that management structures are put in place to ensure sustained action (see *Prosopis juliflora* action plan). Fast-growing annual and perennial weeds such as *Heliotropium curassivicum*, wild tomato (*Solanum* spp.), *Waltheria indica* and *Agremone mexicana* will be more difficult to exclude; however, control should be attempted in a small number of key areas (e.g. Hummock Point and South Gannet Hill) in order to assess on-going management requirements.

Grazing by introduced mammals and invertebrate pests will also be difficult to contain. Fencing has proven to be useful for protecting small numbers of plants from larger mammalian grazers, but is not practical to install or maintain at larger scales. Bell & Boyle [11] assessed various management options for rabbits on Ascension Island, including the feasibility of eradication, and concluded that targeted control using poison baits and traps offered the most tractable solution for reducing their numbers in problematic areas. Regular monitoring of grazing pressure will be needed to assess the magnitude of the threat posed by introduced herbivores and guide management decisions. In terms of invertebrate pests, biological control represents the only viable management option. Indeed, the introduction of the ladybird beetle *Rodolia cardinalis* in 1976 has apparently been effective in controlling a previously serious infestation of cottony cushion scale insect (*Icerya purchasi*) on *E. origanoides* [1,3] and similar biological controls could be explored for other phytophagous pests. A more detailed study of the pest species currently affecting *E. origanoides* and the natural enemies already present on the island would be useful to assess the potential for further biocontrol.

SPECIES ACTION PLAN

PROPOSED ACTION	OUTCOME(S)	TIMEFRAME	PROPOSED START	PRIORITY	LEAD
Policy & Legislation					
Enact planning legislation that mandates open and consultative Environmental Impact Assessments of all major development projects.	1 ordinance enacted	1 year	September 2015	MEDIUM	Head of Conservation (AIG)
Management					
Produce management plans for all nature reserves and sanctuaries, including timetables for the routine control of invasive species, and integrate into the BAP.	6 management plans produced	3 months	March 2015	MEDIUM	Head of Conservation (AIG)
Establish, map and maintain exclusion zones for Mexican thorn, <i>Nicotiana glauca</i> , <i>Casuarina equisetifolia</i> and other invasive shrubs and trees within all coastal and lowland nature reserves.	6 exclusion zones established	3 months	June 2015	HIGH	National Park Warden (AIG)
Maintain cultivated stocks of all <i>Euphorbia origanoides</i> sub-populations as a conservation failsafe and source of material for restoration work.	100 plants in cultivation	1 year then review	September 2014	MEDIUM	Assistant Conservation Officer (AIG)
Attempt to establish at least one new sub-population of <i>Euphorbia origanoides</i> within its former range.	50 individuals re-introduced	1 week	March 2016	MEDIUM	Assistant Conservation Officer (AIG)
Maintain a viable, <i>ex situ</i> seed bank of <i>Euphorbia origanoides</i> seeds on Ascension Island as a conservation failsafe.	100 seeds banked; 1 viability test completed	1 year	January 2015	MEDIUM	Assistant Conservation Officer (AIG)
Research & monitoring					

PROPOSED ACTION	OUTCOME(S)	TIMEFRAME	PROPOSED START	PRIORITY	LEAD
Generate and ground truth a fine-scale, terrestrial habitat map of Ascension Island to enable studies of invasion rates and risks, restoration potential and the distribution, abundance and habitat requirements of native species.	3 GIS layers produced; 1 report/academic paper produced	1 year	April 2015	MEDIUM	Chief Scientist (AIG)
Establish a network of precipitation/temperature monitoring stations to refine habitat classifications and expand the range of baseline meteorological data available for climate change monitoring and research.	6 monitoring stations deployed	1 month	May 2015	MEDIUM	Chief Scientist (AIG)
Produce an endemic plant monitoring manual to clarify the aims of the monitoring programme and document methods for data collection, storage and analysis.	1 report produced	1 month	February 2015	LOW	Assistant Conservation Officer (AIG)
Clean existing endemic plant monitoring data stored in departmental databases and update database structures in line with current monitoring and research needs.	1 database produced	1 year	September 2014	MEDIUM	Assistant Conservation Officer (AIG)
Conduct biannual censuses of <i>Euphorbia orgianoides</i> in known natural populations, timing surveys to coincide with the peaks of the hot and cool seasons,	2 censuses completed	7 months	March 2015	MEDIUM	Assistant Conservation Officer (AIG)
Seek funding and partner support for a project to catalogue Ascension Island's terrestrial invertebrate fauna, including a reassessment of the distribution and status of endemic species and a detailed study of invertebrate pests effecting endemic plants.	1 project initiated	1 year	April 2015	MEDIUM	Head of Conservation (AIG)
Confirm apparent genetic variation between fragmented sub-populations of Ascension spurge and investigate the functional significance of this through common garden experiments.	1 academic paper/thesis produced	3 years	January 2016	MEDIUM	Centre for Ecology & Hydrology

PROPOSED ACTION	OUTCOME(S)	TIMEFRAME	PROPOSED START	PRIORITY	LEAD
Experimentally assess the value of artificially constructed rock shelters as a management technique for increasing local seed bank accumulation around wild-growing Ascension spurge.	1 report/thesis produced	1 year	September 2015	LOW	Chief Scientist (AIG)
Communication & Awareness Raising					
Erect visitor information boards at access points and key features within all nature reserves and national parks, establishing a strong protected areas identity and awareness on Ascension Island.	10 signs installed	6 months	January 2016	MEDIUM	National Park Warden (AIG)
Publish a free guide booklet on Ascension Island's protected areas network to be made available at ports of entry, visitor attractions and other outlets.	500 booklets produced	3 months	June 2015	LOW	Tourism & Media Officers (AIG)
Produce an updated information leaflet on the flora of Ascension Island.	250 leaflets produced	3 months	June 2015	LOW	Tourism & Media Officers (AIG)

Ascension Island BAP: *Euphorbia origanoides*

References

1. Gray A, Robinson PD & Stroud S (2009) Use of the Biological Flora framework in the United Kingdom Overseas Territories: *Euphorbia origanoides* L. *Biological Conservation* **142**, 1754–1766.
2. Duffey E (1964) The terrestrial ecology of Ascension Island. *Journal of Applied Ecology* **1**, 219–251.
3. Cronk QCB (1980) Extinction and survival in the endemic vascular flora of Ascension Island. *Biological Conservation* **17**, 207–219.
4. Lambdon P, Stroud S, Clubbe C, Gray A, Hamilton M, Niissalo M, Pelembe T & Renshaw O (2009) *A plan for the conservation of endemic and native flora on Ascension Island*.
5. Gray A, Stroud S, Lambdon PW, Niissalo M & Renshaw O (2012) *Euphorbia origanoides*. In: *The IUCN Red List of Threatened Species. Version 2014.3*. <www.iucnredlist.org>. [accessed 2015 Feb. 5].
6. Malumphey C, White LF, Hall J, Hodgetts J & Hamilton M (In review) Five invasive mealybug species new for Ascension Island (Hemiptera: Pseudococcidae); with a revised checklist of scale insects for the island.
7. Hughes BJ (2014) *Breeding and population ecology of sooty terns on Ascension Island*. PhD Thesis. University of Birmingham, Birmingham, UK.
8. Gray A (2003) *Ascension Spurge (Euphorbia origanoides L.) study recommendations*. US Air Force.
9. Gray A (2009) *Ascension Spurge Euphorbia origanoides L. climate and viability study: Final Report*. Unpublished report. Centre for Ecology and Hydrology, Edinburgh Research Station.
10. Salafsky N et al. (2008) A Standard Lexicon for Biodiversity Conservation: Unified Classifications of Threats and Actions. *Conservation Biology* **22**, 897–911.
11. Bell E & Boyle D (2008) *Rabbits on Ascension Island: Assessment of the potential for eradicating rabbits from Ascension and developing a rabbit control strategy for immediate implementation*. Wildlife Management International Limited.