

# Ascension Island Biodiversity Action Plan

## GREEN TURTLE



### SUMMARY

**Taxonomy:** Kingdom: Animalia; Phylum: Chordata; Class: Reptilia; Order: Testudines; Family: Cheloniidae; Species: *Chelonia mydas*

**Nativeness:** Native, breeding

**Description:** Large, highly migratory marine reptile up to 1.5 metres in length and 300 kg in weight. Migrates from coastal feeding areas along the Brazilian continental shelf to nest on Ascension Island's sandy beaches. Adults are predominantly herbivorous with a diet composed largely of seaweed and seagrass.

**IUCN Red List status:** Endangered 

**Local trend:** Increasing 

**Threats:** Principle threats are climate change and fisheries bycatch (particularly in coastal feeding areas), with secondary threats from invasive rodents and nesting habitat degradation by invasive weeds, light pollution and erosion.

Distribution	
<b>Global</b>	
Occurs circum-globally in tropical and subtropical seas. Nests in over 80 countries worldwide, with major Atlantic breeding populations located in Costa Rica, Ascension Island (UK), Poilao (Guinea-Bissau), Florida, Suriname, Mexico, Trindade (Brazil), Bioko (Equatorial Guinea) and Aves Island (Venezuela) [1].	
<b>Local</b>	
<p><b>Nesting:</b> Green turtle nesting at Ascension Island occurs between December and June on numerous sandy beaches and coves located along the western, northern and north-eastern coastlines (Fig. 1). Approximately 75% of nesting occurs on three primary beaches: Long Beach, Pan Am Beach and North East Bay (Fig. 1) [2]. Sites of secondary importance include Deadman’s Beach and Clarke’s Beach. During the mating season (Nov – Mar), large aggregations of mating turtles can be found in shallow, inshore waters, particularly in bays adjacent to the primary nesting beaches [3]. Females reside in shallow sub-littoral habitats during the intervals between nesting events, typically resting at depths shallower than 20 m [4].</p> <p><b>Foraging:</b> Adult green turtles nesting at Ascension Island forage along the Brazilian coastal shelf from Rio de Janeiro to Fortaleza [5,6]. Post-nesting migrations are biphasal, comprising of a westerly open ocean crossing making landfall around the ‘bulge’ of Brazil, followed by a period of extended coastal travel to their final destination [6]. Juvenile green turtles originating from Ascension are even more widely distributed, having been identified at foraging grounds along a 6000 km stretch of South American coastline from northern Argentina to the Caribbean ([7] and refs therein).</p>	<p><b>Figure 1.</b> Distribution of green turtle nesting during the last full survey in 2011-2012 [2]. Symbol sizes are scaled according to total nesting activity.</p>

3. Status				
GLOBAL	Population estimate:	Unknown	IUCN status:	Endangered
Global population size of the green turtle has not been formally assessed. Based on an evaluation of 34 index sites for which data are available [8] estimated a 48-67 % decline over three generations, justifying its classification as endangered on the IUCN Red List. However, many populations are currently growing rapidly (e.g. [9]) and its endangered status has been strongly questioned [10].				
LOCAL	Population estimate:	ca. 15,000 adult females	Local trend:	Increasing
Following the discovery of Ascension Island in the 16th century, the green turtle population was subjected to centuries of intensive exploitation for meat, resulting in a severe depletion of the population [10,11]. Harvesting had largely ceased by the 1940s [10], and, since monitoring began in the 1970s, estimated annual numbers of green turtle nests have increased 6-fold from approximately 3750 to more than 23,000 [2]. The status of the population is therefore currently considered to be favourable.				

4. Ecology
Habitat & diet
Green turtles occupy different habitats during different life stages. Adults are predominantly herbivorous and undertake cyclical, long-distance migrations between coastal feeding habitats and nesting sites, which are often located on oceanic islands. Hatchlings and small juveniles are epipelagic omnivores and are thought to associate with

floating vegetation and other debris entrained in ocean currents [12].
<b>Reproduction &amp; life history</b>
Green turtles nest on sandy marine beaches. Females lay an average of 6 clutches of 120 eggs within a nesting season and breed at intervals of 3-4 years [13,14]. Sex is determined by nest temperature during the middle third of incubation, with a pivotal temperature (giving an equal sex ratio) of approximately 29 C [15]. Hatchlings emerge 45 – 60 days after nesting, normally at night, and disperse rapidly into the open ocean. Juveniles join adults in coastal feeding grounds at a carapace length of approximately 30 cm and are estimated to reach maturity at 17 – 35 years of age [16].
<b>Taxonomy &amp; population structure</b>
Green turtles tend to breed at their natal nesting sites resulting in reproductive isolation among nesting populations [17]. The Ascension Island population is genetically distinct from other Atlantic populations [17] and there is evidence of local divergence between turtles nesting at east coast and west coast beaches [18], although local genetic structure appears to be weak [19,20].

<b>4. Threats*</b>		
<b>11.1 Climate change &amp; severe weather: habitat shifting and alteration</b>	<b>Impact:</b>	<b>MEDIUM</b>
Loss of suitable nesting habitat as a result of climate change-induced sea level rise poses a significant threat to marine turtles worldwide, particularly in areas where coastal geomorphology and development curtail the landward migration of beaches [21]. Sea level at Ascension Island is estimated to have risen by 7 cm since 1955 [22] and this trend is projected to accelerate during the first half of the 21st century [23]. Cliffs, roads and other coastal land use represent barriers to beach migration in some areas and could result in ‘coastal squeeze’ if not addressed.		
<b>11.3 Climate change &amp; severe weather: temperature extremes</b>	<b>Impact:</b>	<b>MEDIUM</b>
Rising temperatures on nesting beaches are predicted to become a major threat to marine turtles during the 21st century, skewing sex ratios (marine turtles have temperature-dependent sex determination) and reducing the hatching success of eggs [21]. The primary sex ratio at Ascension Island is already estimated to be 75% female [15] and incubation temperatures on some beaches are close to the upper limits for embryonic development during much of the nesting season [18]. Incubation temperatures have increased by an estimated 0.5 C over the past 100 years [24], and this trend is projected to accelerate during the 21st century [23].		
<b>5.4.4 Fishing &amp; harvesting aquatic resources (unintentional effects, large scale)</b>	<b>Impact:</b>	<b>MEDIUM</b>
Incidental capture in fisheries constitutes a major source of mortality for marine turtles worldwide and may affect green turtles both in their coastal foraging habitats and during oceanic migrations. Reproductive migrations between Brazil and Ascension Island traverse an area of intensive pelagic long-lining for tuna and billfish, which is estimated to kill thousands of marine turtles annually across the Atlantic region [25]. Green turtles account for a relatively small proportion of this total, with most studies in the southern and central Atlantic reporting a by-catch rate of 0-13 individuals per million hooks [26–28]. The herbivorous diet and predominantly coastal distribution of adult green turtles probably explain this reduced susceptibility, although specific data from along oceanic migration routes where risks may be elevated is lacking. In contrast, by-catch rates of green turtles in coastal trawl, weir and net fisheries in the south west Atlantic are substantial [28,29] and may impact both adults and juvenile stages originating from Ascension Island.		
<b>8.1.2 Invasive non-native/alien species/diseases (named species)</b>	<b>Impact:</b>	<b>MEDIUM</b>
Invasive ship rats ( <i>Rattus rattus</i> ) predate green turtle hatchlings and are locally abundant around some nesting beaches, most notably North East Bay where significant levels of hatchling predation has been reported [30]. Their numbers appear to be increasing following a successful campaign to eradicate feral cats between 2001 and 2004 [31]. Several species of invasive, salt-tolerant plants, including <i>Prosopis juliflora</i> , <i>Nicotiana glauca</i> , <i>Argemone mexicana</i> , <i>Heliotropium curassavicum</i> , <i>Waltheria indica</i> and <i>Chenopodium murale</i> , are also increasingly encroaching on nesting beaches and threaten to reduce the extent of suitable nesting habitat. Dense growths of weeds and shrubs can directly obstruct nesting activity and alter substrate composition, consolidating the ground and contributing to soil formation which further facilitates the invasion process. The tree <i>Casuarina equisetifolia</i> is		

also rapidly expanding its range in the east of the Island and is a notoriously aggressive invader of sand beaches [32,33]. It has not yet reached the turtle nesting beaches of the east coast and must be prevented from doing so.		
<b>9.6.1 Light pollution</b>	<b>Impact:</b>	LOW
Anthropogenic lighting disrupts marine turtle nesting and disorients hatchlings, preventing them from reaching the sea. Low-level light pollution is responsible for some hatchling mortality on nesting beaches close to Georgetown and the US Air Force Base, but impacts appear to be limited. Only the beach area directly adjacent to the Pierhead is exposed to continuous, potentially disruptive levels of artificial lighting and could benefit from mitigation measures. Beach hut parties, night time refuelling operations at the Catherine Point depot and vessels moored in the Clarence Bay anchorage are also intermittent sources of light pollution. Lights from the latter are known to attract hatchlings and may expose them to increased predation by large, pelagic fish that school underneath.		
<b>9.2.1 Industrial &amp; military effluents (oil spills)</b>	<b>Impact:</b>	LOW
Although unlikely, a major petrochemical spill at Ascension Island could have disastrous impacts on the green turtle nesting population, especially given the proximity of the tanker anchorage, ship-to-shore refuelling operations, storage silos and pipelines to some of the Island's most important nesting habitat.		
<b>6.1 Recreational activities</b>	<b>Impact:</b>	LOW
Viewing nesting turtles is one of the most popular visitor experiences on Ascension Island. While this presents opportunities for awareness-raising, some unintentional disturbance of nesting females inevitably occurs, particularly on Long Beach which is the focus for most 'turtle tourism'. Impacts are currently limited by low visitor numbers, but any plans to increase the tourism footprint on Ascension Island should consider measures to minimise disruption to nesting turtles.		
<b>1. Residential &amp; commercial development</b>	<b>Impact:</b>	NEGLIGIBLE
Coastal development is currently limited around most nesting beaches, although the construction of roads, pipelines and parking areas have encroached on marginal nesting habitat in some areas. Greater conflicts between coastal land use and the preservation of turtle nesting habitat may arise in future if predicted sea level rise eventuates. All beachfront land is either publicly owned or under military concession making development control comparatively straightforward given the correct regulatory framework.		
<b>9.1.2 Domestic &amp; urban waste water (run-off)</b>	<b>Impact:</b>	NEGLIGIBLE
Localised erosion caused by rapid storm water run-off from developed areas has degraded some areas of nesting habitat, particularly the southern end of Long Beach which currently supports limited nesting activity.		
<b>3.2 Mining &amp; quarrying</b>	<b>Impact:</b>	NEGLIGIBLE
Mining of beach sand for construction was once commonplace on Ascension Island and has resulted in a significant and lasting reduction in the extent of suitable nesting habitat [34–36]. The practice has been subject to an unofficial moratorium since the mid-1990s, although permission to remove small volumes of sand has still occasionally been granted. An environmental impact assessment commissioned in 2004 suggested that 78% of nesting beaches were showing some evidence of erosion and recommended that no further sand extraction should be permitted [36].		
*Threats are classified and scored according to the <a href="#">IUCN-CMP Unified Classification of Direct Threats</a> [37]		

Relevant policies and legislation	
International	
The green turtle is listed under <a href="#">Appendix I of CITES</a> (commits parties to adopt legislation prohibiting the import of export of listed species without license from local scientific and management authorities). Although provisions for implementing CITES locally are contained within the Endangered Species (Ascension) Control Ordinance 1967 this legislation has been highlighted as inadequate and is in the process of being updated.	
The green turtle is listed under <a href="#">Appendix I of the Convention on Migratory Species</a> (commits parties to conserving and restoring the habitats of listed species [Article III, para. 4a], minimising activities that impede their migration [Article III, paragraph 4b] and prohibiting the taking of those species [Article III, para. 5]).	

**Local**

Green turtles are protected under the [Wildlife Protection Ordinance 2013](#), which prohibits the killing, capture or taking of turtles or their eggs on Ascension Island without license.

The major green turtle nesting beaches of Long Beach, Pan Am Beach and North East Bay as well as their hinterlands and adjacent near shore waters are designated as Nature Reserves under the [National Protected Areas Order 2014](#) and [National Protected Areas Regulations 2014](#). The regulations restrict all forms of development within beach reserves and contain provisions to control pets, open fires, vehicles and motor craft, beach hut use and other forms of recreation.

**6. Management notes**

Green turtles and their nesting habitat are already well-protected locally and the major threats to the population that remain, including climate change and fisheries by-catch on the South American feeding grounds, are largely beyond local control. Nevertheless, there may be steps that can be taken locally to improve population resilience. The rapidly increasing rat population on the Island is the most immediate concern and if left unmanaged may significantly depress green turtle productivity as a result of predation on hatchlings. The deployment of poison baits at North East Bay appears to have been successful in reducing rat densities and hatchling predation at this site and should be implemented at all major nesting beaches, along with improved monitoring to assess the efficacy of control measures. Various other anthropogenic factors including invasive weeds, light pollution, urban runoff and human disturbance also currently affect nesting areas and, although individually limited in severity, may have an additive effect on reducing the overall quality of nesting habitat and should be addressed when possible. Formal habitat management plans for major nesting beaches are needed to assign responsibilities and ensure continuity of action, particularly with regards to invasive species control.

Despite the favourable status of the population, annual monitoring of green turtle nesting activity should continue over the short to medium term, particularly given its importance as a regional indicator population for the south Atlantic. Currently, nesting activity on the three main beaches is monitored on two consecutive days every 1 – 2 weeks and there is no reason to increase this effort. Although this work is labour intensive, the continuation of an internship programme initiated in 2014 will hopefully minimise the impact of turtle monitoring on other more urgent conservation commitments. A full census on all nesting beaches should be conducted periodically (5 – 10 years) to calibrate population estimates derived from index sites (see Weber et al. 2014) and regular monitoring of hatching success and nest temperatures would be valuable given the potential threats from climate change. Indeed, rising global temperatures and sea levels present the greatest long-term threat to the survival of the Ascension Island green turtle population and may be difficult or impossible to manage locally. Nonetheless, research into the likely impacts of climate change on the population is underway and will hopefully provide insights into possible mitigation measures, such as combatting ‘coastal squeeze’ by providing room for the natural migration of nesting beaches.

**SPECIES ACTION PLAN**

PROPOSED ACTION	OUTCOME(S)	TIMEFRAME	PROPOSED START	PRIORITY	LEAD*
Policy & Legislation					
Enact planning legislation that mandates open and consultative Environmental Impact Assessments of all major development projects.	1 ordinance enacted	1 year	September 2015	MEDIUM	Head of Conservation (AIG)
Update existing legislation underpinning CITES to comply with the basic legislative requirements of the Convention.	1 ordinance enacted	1 year	July 2014	LOW	Solicitor General (AIG)
Develop an implementation plan for reforming Ascension Island's fishing policies and marine protection legislation, taking into account the recommendations of previous consultancy reports and the outcome of deliberations on the future of the Territory's commercial offshore fishery.	1 policy document;	1 year	March 2015	HIGH	Director of Fisheries (AIG)
Review existing petrochemical spill response plans associated with ship-to-shore refuelling activities and petroleum storage, ensuring that financial and operational responsibilities are clearly defined and copies are lodged with all stakeholders.	1 memorandum of understanding	2 months	June 2015	MEDIUM	Director of Operations (AIG)
Enact a legislative ban on sand extraction from beaches and offshore deposits.	1 ordinance enacted	6 months	September 2015	MEDIUM	Head of Conservation (AIG)
Management					
Produce management plans for all nature reserves and sanctuaries, including timetables for the routine control of invasive species, and integrate into the BAP.	6 management plans produced	1 year	March 2015	MEDIUM	Head of Conservation (AIG)

PROPOSED ACTION	OUTCOME(S)	TIMEFRAME	PROPOSED START	PRIORITY	LEAD*
Recruit, train and equip two permanent, core-funded rangers to assist with the implementation of protected areas management plans, in particular the control of invasive species encroaching on key wildlife habitat and amenity areas.	2 staff recruited	2 months	April 2015	MEDIUM	Head of Conservation (AIG)
Establish, map and maintain exclusion zones for <i>Prosopis juliflora</i> , <i>Nicotiana glauca</i> , <i>Casuarina equisetifolia</i> and other invasive drought-tolerant shrubs and trees within all coastal reserves.	6 exclusion zones cleared and mapped	6 months	June 2015	HIGH	National Park Warden (AIG)
Identify and source sustainable alternatives to beach sand as a line-marking medium for public sports fields.	1 product sourced	3 months	April 2015	LOW	Director of Operations (AIG)
Restore green turtle nesting habitat at Long Beach degraded by erosion, sand extraction and coastal land use through drainage improvements, beach re-nourishment and/or reclamation of parking areas and other degraded sites.	0.25 hectares restored	1 year	April 2015	MEDIUM	Head of Conservation (AIG)
Install permanent rodent control lines at all primary green turtle nesting beaches supporting more than 5 % of total annual nesting activity.	40 stations deployed	1 month	November 2015	MEDIUM	Environmental Health (AIG)
Research & monitoring					
Carry out a comprehensive assessment of anthropogenic light pollution and hatchling disorientation on all major green turtle nesting beaches, including recommendations for remedial action and standards for future beachfront lighting.	1 report produced	3 months	April 2016	MEDIUM	Chief Scientist (AIG)
Organise an annual marine turtle internship programme to provide career relevant training for recent graduates and help deliver the objectives of the green turtle action plan.	3 interns recruited	2 months	September 2015	MEDIUM	Head of Conservation (AIG)

PROPOSED ACTION	OUTCOME(S)	TIMEFRAME	PROPOSED START	PRIORITY	LEAD*
Estimate annual green turtle nesting activity on Long Beach, Pan Am Beach and North East Bay through routine counts of tracks and nests between December and June.	1 report produced	7 months	December 2014	MEDIUM	Marine Turtle Interns (AIG)
Carry out a complete survey of green turtle nesting activity on all of Ascension Island's beaches in accordance with previously published methods.	1 report produced	7 months	December 2016	LOW	Marine Turtle Interns (AIG)
Assess the abundance of black rats at North East Bay following the deployment of bait stations in response to high predation rates on green hatchlings recorded in 2013.	1 report produced	5 days	May 2015	MEDIUM	Marine Turtle Interns (AIG)
Initiate routine indexing of rat abundance at Long Beach, Pan Am Beach and North East Bay during the peak of green turtle hatchling emergence.	30 stations deployed; 1 survey completed	2 weeks	May 2016	MEDIUM	Marine Turtle Interns (AIG)
Monitor the incubation temperature and hatching success of a spatially and temporally representative sample of green turtle clutches on Long Beach, Pan Am Beach and North East Bay in each nesting season	60 nests monitored; 1 report produced	7 months	January 2015	MEDIUM	Marine Turtle Interns (AIG)
Experimentally determine the pivotal sex determining temperature of green turtles at Ascension Island to parameterise climate change sensitivity models	1 academic paper/thesis produced	6 months	January 2015	MEDIUM	University of Exeter
Produce accurate elevation maps of primary green turtle nesting beaches and model potential losses from predicted sea level rise, producing recommendations for possible mitigation strategies.	1 academic paper/thesis produced	6 months	January 2016	MEDIUM	University of Exeter
Develop a robust and practical system for monitoring changes in green turtle nesting beach profiles and extents.	1 report produced	6 months	January 2016	MEDIUM	University of Exeter

PROPOSED ACTION	OUTCOME(S)	TIMEFRAME	PROPOSED START	PRIORITY	LEAD*
Develop collaborations to collate existing data on contemporary sea level rise at Ascension Island and implement systems for long-term sea level monitoring and reporting.	1 database created; 1 action plan produced	6 months	June 2016	LOW	Chief Scientist (AIG)
Work with the international research community to consolidate existing leatherback turtle and green turtle tracking data collected within Ascension Island's EEZ and integrate into GIS as a basis for marine spatial planning.	2 GIS layers created	6 months	May 2016	LOW	Chief Scientist (AIG)
Communication & Awareness Raising					
Erect visitor information boards at access points and key features within all nature reserves and national parks, establishing a strong protected areas identity and awareness on Ascension Island.	10 signs installed	6 months	January 2016	MEDIUM	National Park Warden (AIG)
Publish a free guide booklet on Ascension Island's protected areas network to be made available at ports of entry, visitor attractions and other outlets.	500 booklets produced	3 months	June 2015	LOW	Tourism & Media Officers (AIG)
Celebrate World Ocean's Day as a focus for awareness raising activities associated with Ascension Island's marine environment.	1 event organised	1 day	June 2015	LOW	Head of Conservation (AIG)
Initiate an education campaign on the importance of not discarding plastics and fishing tackle at sea, including the installation of signage and tackle bins at known fishing hotspots.	3 signs/bins installed	3 months	May 2015	LOW	Head of Conservation (AIG)
Produce an updated visitor information leaflet on the biology and conservation of marine turtles at Ascension Island.	500 leaflets produced	1 month	March 2015	LOW	Tourism & Media Officers (AIG)
Produce and widely publicise a code of conduct for turtle watchers and beach hut users during turtle nesting season.	Code of conduct displayed at all beach huts and major nesting beaches	3 months	October 2015	MEDIUM	Tourism & Media Officers (AIG)

PROPOSED ACTION	OUTCOME(S)	TIMEFRAME	PROPOSED START	PRIORITY	LEAD*
Upload all existing green turtle monitoring data from Ascension Island into the State of the World's Turtles (SWOT) online repository	1 dataset uploaded	2 days	July 2015	LOW	Chief Scientist (AIG)
Organise regular guided 'turtle tours' for island residents and visitors to view nesting turtles and learn about their biology and conservation	20 tours organised	5 months	January 2015	MEDIUM	Marine Turtle Interns (AIG)
Arrange a public 'beach clean' event to encourage members of the local community to participate in the preservation of the Island's sea turtle nesting habitat and recreation spaces	1 event organised	1 day	October 2015	MEDIUM	Conservation Assistant (AIG)
Convert the former Saints' Beach Hut on Long Beach into a visitor information centre and learning space, providing a focal point for awareness raising activities associated with marine turtles and the wider marine environment	1 visitor centre created	1 year	January 2015	LOW	Head of Conservation (AIG)

## References

1. De Padua Almeida A, Moreira LM, Bruno SC, Thomé JCA, Martins AS, Bolten AB & Bjorndal KA (2011) Green turtle nesting on Trindade Island, Brazil: abundance, trends, and biometrics. *Endangered Species Research* **14**, 193–201.
2. Weber SB, Weber N, Ellick J, Avery A, Godley BJ, Sim J, Williams N & Broderick AC (2014) Recovery of the South Atlantic's largest green turtle nesting population. *Biodiversity and Conservation* **23**, 3005–3018.
3. Godley BJ, Broderick AC, Frauenstein R, Glen F & Hays GC (2002) Reproductive seasonality and sexual dimorphism in green turtles. *Marine Ecology Progress Series* **226**, 125–133.
4. Hays GC, Adams CR, Broderick AC, Godley BJ, Lucas DJ, Metcalfe JD & Prior AA (2000) The diving behaviour of green turtles at Ascension Island. *Animal Behaviour* **59**, 577–586.
5. Carr A (1975) The Ascension Island green turtle colony. *Copeia* **1975**, 547–555.
6. Hays G, Broderick AC, Godley BJ, Lovell P, Martin C, McConnell BJ & Richardson S (2002) Biphasal long-distance migration in green turtles. *Animal Behaviour* **64**, 895–898.
7. Naro-Maciel E, Bondioli ACV, Martin M, de Padua Almeida A, Baptistotte C, Bellini C, Marcovaldi MA, Santos AJB & Amato G (2012) The Interplay of Homing and Dispersal in Green Turtles: A Focus on the Southwestern Atlantic. *Journal of Heredity* **103**, 792–805.
8. Seminoff JA (2004) *Chelonia mydas*. In: *IUCN Red List of Threatened Species. Version 2014.3*. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 04 February 2015.
9. Chaloupka M, Bjorndal KA, Balazs GH, Bolten AB, Ehrhart LM, Limpus CJ, Suganuma H, Troëng S & Yamaguchi M (2008) Encouraging outlook for recovery of a once severely exploited marine megaherbivore. *Global Ecology and Biogeography* **17**, 297–304.
10. Broderick AC, Frauenstein R, Glen F, Hays GC, Jackson AL, Pelembe T, Ruxton GD & Godley BJ (2006) Are green turtles globally endangered? *Global Ecology and Biogeography* **15**, 21–26.
11. Huxley RC (1999) Historical overview of marine turtle exploitation, Ascension Island, South Atlantic. *Marine Turtle Newsletter* **84**, 7–9.
12. Witherington B, Hirama S & Hardy R (2012) Young sea turtles of the pelagic Sargassum-dominated drift community: habitat use, population density, and threats. *Marine Ecology Progress Series* **463**, 1–22.
13. Mortimer JA & Carr A (1987) Reproduction and migrations of the Ascension Island green turtle (*Chelonia mydas*). *Copeia* **1987**, 103–113.
14. Weber N, Weber SB, Godley BJ, Ellick J, Witt M & Broderick AC (2013) Telemetry as a tool for improving estimates of marine turtle abundance. *Biological Conservation* **167**, 90–96.
15. Godley BJ, Broderick AC, Glen F & Hays GC (2002) Temperature-dependent sex determination of Ascension Island green turtles. *Marine Ecology Progress Series* **226**, 115–124.
16. Hirth HF (1997) *Synopsis of biological data on the green turtle Chelonia mydas*. US Fish and Wildlife Service.
17. Bowen BW, Meylan AB, Ross TP & Limpus CT (1992) Global population structure and natural history of the green turtle (*Chelonia mydas*) in terms of matriarchal phylogeny. *Evolution* **46**, 865–881.
18. Weber SB, Broderick AC, Groothuis TGG, Ellick J, Godley BJ & Blount JD (2012) Fine-scale thermal adaptation in a green turtle nesting population. *Proceedings of the Royal Society B: Biological Sciences* **279**, 1077–1084.
19. Formia A, Broderick AC, Glen F, Godley BJ, Hays GC & Bruford MW (2007) Genetic composition of the Ascension

- Island green turtle rookery based on mitochondrial DNA: implications for sampling and diversity. *Endangered Species Research* **3**, 145–158.
20. Lee PLM, Luschi P & Hays GC (2007) Detecting female precise natal philopatry in green turtles using assignment methods. *Molecular Ecology* **16**, 61–74.
  21. Hawkes L, Broderick A, Godfrey M & Godley B (2009) Climate change and marine turtles. *Endangered Species Research* **7**, 137–154.
  22. Woodworth P et al. (2012) Sea level changes at Ascension Island in the last half century. *African Journal of Marine Science* **34**, 443–452.
  23. IPCC (2013) Summary for Policymakers. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds. T. F. Stocker D. Qin G.-K. Plattner M. Tignor S. K. Allen J. Boschung A. Nauels Y. Xia V. Bex & P. M. Midgley). Cambridge University Press, Cambridge, UK. pp. 1–30.
  24. Hays GC, Broderick AC, Glen F & Godley BJ (2003) Climate change and sea turtles: a 150-year reconstruction of incubation temperatures at a major marine turtle rookery. *Global Change Biology* **9**, 642–646.
  25. Angel A, Nel R, Wanless RM, Mellet B, Harris L & Wilson I (2014) Ecological risk assessment of sea turtles to tuna fishing in the ICCAT region. *Collective Volume of Scientific Papers ICCAT* **70**, 2226–2259.
  26. Sales G, Giffoni BB & Barata PCR (2008) Incidental catch of sea turtles by the Brazilian pelagic longline fishery. *Journal of the Marine Biological Association UK* **88**, 853–864.
  27. Coelho R, Fernandez-Carvalho J & Santos MN (2013) A review of fisheries within the ICCAT convention area that interact with sea turtles. *Collective Volume of Scientific Papers ICCAT* **69**, 1788–1827.
  28. Wallace BP, Kot CY, DiMatteo AD, Lee T, Crowder LB & Lewison RL (2013) Impacts of fisheries bycatch on marine turtle populations worldwide: toward conservation and research priorities. *Ecosphere* **4**, art40.
  29. Gallo BMG, Macedo S, Giffoni B de B, Becker JH & Barata PCR (2006) Sea turtle conservation in Ubatuba, southeastern Brazil, a feeding area with incidental capture in coastal fisheries. *Chelonian Conservation and Biology* **5**, 93–101.
  30. Dawson E (2013) *The diet and distribution of invasive ship rats (Rattus rattus) on Ascension Island*. MSc Thesis. University of Exeter, Exeter, UK.
  31. Hughes BJ (2014) *Breeding and population ecology of sooty terns on Ascension Island*. PhD Thesis. University of Birmingham, Birmingham, UK.
  32. Global Invasive Species Database (2015) *Casuarina equisetifolia*. Available from <http://www.issg.org/database/species/ecology.asp?si=365&fr=1&sts=sss&lang=EN> [Accessed 04 February 2015].
  33. Awale D & Phillott AD (2014) A review of the adverse effects of *Casuarina* spp. on coastal ecosystems and sea turtle nesting beaches. *Indian Ocean Turtle Newsletter* **19**, 15–19.
  34. Fletemeyer J (1986) Guest Editorial: Ascension Island Turtles Still in Trouble. *Marine Turtle Newsletter* **37**, 11.
  35. Mortimer JA (1992) *An evaluation of the status of the Ascension Island green turtles and their nesting habitat in 1992*. Unpublished report. University of Florida / Caribbean Conservation Corporation.
  36. Cambers G (2004) *Environmental Impact Assessment: Beach sand extraction on Ascension Island, South Atlantic*. University of Puerto Rico.

37. Salafsky N et al. (2008) A Standard Lexicon for Biodiversity Conservation: Unified Classifications of Threats and Actions. *Conservation Biology* **22**, 897–911.